

THE BISHOP-PHELPS-BOLLOBÁS MODULUS FOR FUNCTIONALS ON CLASSICAL BANACH SPACES

F. J. GARCÍA-PACHECO* and S. MORENO-PULIDO

Communicated by P. Aiena

ABSTRACT. In this manuscript we compute the Bishop-Phelps-Bollobás modulus for functionals in classical Banach spaces, such as Hilbert spaces, spaces of continuous functions c_0 and ℓ_1 .

REFERENCES

1. R. M. Aron, Y. S. Choi, S. K. Kim, H. J. Lee, and M. Martín, *The Bishop-Phelps-Bollobás of Lindenstrauss properties A and B*, Trans. Amer. Math. Soc. **367** (2015), no. 9, 6085–6101.
2. F. J. García-Pacheco and S. Moreno-Pulido, *The Bishop-Phelps-Bollobás modulus for operators*, Preprint.
3. F. J. García-Pacheco and J. R. Hill, *Geometric characterizations of Hilbert spaces*, Canad. Math. Bull. **59** (2016), no. 4, 769–775.
4. M. Chica, V. Kadets, M. Martín, S. Moreno-Pulido, and F. Rambla-Barreno, *Bishop-Phelps-Bollobás moduli of a Banach space*, J. Math. Anal. Appl. **412** (2014), no. 2, 697–719.

DEPARTMENT OF MATHEMATICS, COLLEGE OF ENGINEERING, UNIVERSITY OF CADIZ, PUERTO REAL 11519, SPAIN.

E-mail address: garcia.pacheco@uca.es

E-mail address: soledad.moreno@uca.es

Copyright 2018 by the Tusi Mathematical Research Group.

Date: Received: Dec. 24, 2017; Accepted: Mar. 11, 2018.

*Corresponding author

\diamond Advance publication – final volume, issue, and page numbers to be assigned.

2000 *Mathematics Subject Classification.* Primary 47A05; Secondary 46B20.

Key words and phrases. Bishop-Phelps-Bollobás, modulus, classical Banach spaces.