

TITLE OF PAPER

FIRST AUTHOR¹, SECOND AUTHOR¹, and THIRD AUTHOR^{2*}

ABSTRACT. AOT is now included in the Emerging Sources Citation Index and has been accepted for inclusion in Scopus. The journal is an author-prepared journal, which means that authors are responsible for the proper formatting of manuscripts by using the style file of the journal.

1. INTRODUCTION AND PRELIMINARIES

The Advances in Operator Theory (AOT) is NOT an open access journal. It is a peer-reviewed quarterly electronic journal. It is hosted by Project Euclid, an online platform designed specifically for mathematicians. Project Euclid is a joint venture of Duke University Press and Cornell University. The publisher is the Tusi Math. Research Group. By submitting a manuscript, the author(s) agree that the copyright for the article is transferred to the publisher, if and when, the paper is accepted for publication.

Here you should state the introduction, preliminaries and your notation. Authors are required to state clearly the contribution of the paper and its significance in the introduction. There should be some survey of relevant literature.

1.1. Instructions for author(s). Manuscripts should be typeset in English with double spacing by using AMS-LaTeX. The authors are encouraged to use the journal style file that has been developed for LaTeX2e standard and can be found at the website of the journal.

While you are preparing your paper, please take care of the following:

- (1) Abstract: 200 words or less with no reference number therein.
- (2) MSC2010: Primary only one item; and Secondary at least one item (We need at least one in MSC43, MSC46 or MSC47).
- (3) Keywords: At least 3 items and at most 5 items.
- (4) Authors: Full names, mailing addresses and emails of all authors.

Copyright 2016 by the Tusi Mathematical Research Group.

Date: Received: xxxxxx; Accepted: zzzzzz.

*Corresponding author

◇ Advance publication – final volume, issue, and page numbers to be assigned.

2010 *Mathematics Subject Classification.* Primary 46L05; Secondary 44B20, 46C05.

Key words and phrases. convexity, dual space, C^* -algebra, Hahn–Banach theorem.

- (5) Each Theorem, Proposition, Corollary, Lemma, Definition, Example, etc should be typeset in its respective environment such as $\backslash\begin{theorem}...\backslash\end{theorem}$ and so on.
- (6) Margins: A long formula should be broken into two or more lines. Empty spaces in the text should be removed.
- (7) Tags (Formula Numbers): Equations and numbered items referred to in the text must be labeled by $\backslash\label{A}$. References to them must be typeset by using $\backslash\eqref{A}$. Remove unused tags. Manual numbering of equations or sections must be avoided.
- (8) Acknowledgments: At the end of paper but preceding to References.
- (9) References: Use $\backslash\cite{MM}$ to refer to the specific book/paper [3] (with $\backslash\textit{bibitem}{MM}$) in the text. Remove unused references. References should be listed in the alphabetical order according to the surnames of the first author at the end of the paper and should be cited in the text as, e.g., [2] or [2, Theorem 4.2], etc.
- (10) Abbreviations: Abbreviations of titles of periodicals/books should be given by using Math. Reviews, see Abbreviations of names of serials or MR lookup.

2. MAIN RESULTS

The following is an example of a definition.

Definition 2.1. Let \mathcal{X} be a real or complex linear space. A mapping $\|\cdot\| : \mathcal{X} \rightarrow [0, \infty)$ is called a 2-norm on \mathcal{X} if it satisfies the following conditions:

- (1) $\|x\| = 0 \Leftrightarrow x = 0$,
- (2) $\|\lambda x\| = \|\lambda\| \|x\|$ for all $x \in \mathcal{X}$ and all scalar λ ,
- (3) $\|x + y\|^2 \leq 2(\|x\|^2 + \|y\|^2)$ for all $x, y \in \mathcal{X}$.

Here is an example of a table.

TABLE 1.

1	2	3
$f(x)$	$g(x)$	$h(x)$
a	b	c

This is an example of a matrix

$$\begin{bmatrix} 1 & -2 \\ 3 & 5 \end{bmatrix}$$

The following is an example of an example.

Example 2.2. Let $\theta : \mathcal{A} \rightarrow \mathcal{A}$ be a homomorphism. Define $\varphi : \mathcal{A} \rightarrow \mathcal{A}$ by $\varphi(a) = a_0\theta(a)$. Then we have

$$\begin{aligned}\varphi(a_1 \dots a_n) &= a_0\theta(a_1 \dots a_n) \\ &= \varphi(a_1) \dots \varphi(a_n).\end{aligned}\tag{2.1}$$

Hence φ is an n -homomorphism.

The following is an example of a theorem and a proof. Please note how to refer to a formula.

Theorem 2.3. *If \mathbf{B} is an open ball of a real inner product space \mathcal{X} of dimension greater than 1, \mathcal{Y} is a real sequentially complete linear topological space, and $f : \mathbf{B} \setminus \{0\} \rightarrow \mathcal{Y}$ is orthogonally generalized Jensen mapping with parameters $s = t > \frac{1}{\sqrt{2}}r$, then there exist additive mappings $T : \mathcal{X} \rightarrow \mathcal{Y}$ and $b : \mathbb{R}_+ \rightarrow \mathcal{Y}$ such that $f(x) = T(x) + b(\|x\|^2)$ for all $x \in \mathbf{B} \setminus \{0\}$.*

Proof. First note that if f is a generalized Jensen mapping with parameters $t = s \geq r$, then

$$\begin{aligned}f(\lambda(x + y)) &= \lambda f(x) + \lambda f(y) \\ &\leq \lambda(f(x) + f(y)) \\ &= f(x) + f(y)\end{aligned}\tag{2.2}$$

for some $\lambda \geq 1$ and all $x, y \in \mathbf{B} \setminus \{0\}$ such that $x \perp y$. Now the result can be deduced from (2.2). \square

The following is an example of a remark.

Remark 2.4. One can easily conclude that g is continuous by using Theorem 2.3.

Again, note how we refer to Theorem 2.3 and formula (2.1).

Acknowledgments. Acknowledgments may be placed at the end of the text, immediately preceding the references.

REFERENCES

1. U. Haagerup, *Solution of the similarity problem for cyclic representations of C^* -algebras*, Ann. of Math. (2) **118** (1983), no. 2, 215–240.
2. G. J. Murphy, *C^* -Algebras and Operator Theory*, Academic Press, Boston, 1990.
3. M. Mirzavaziri and M. S. Moslehian, *Automatic continuity of σ -derivations in C^* -algebras*, Proc. Amer. Math. Soc. **134** (2006), no. 11, 3319–3327.
4. M. S. Moslehian, *Ky Fan inequalities*, Linear Multilinear Algebra, arXiv:1108.1467v2 (to appear).
5. Th. M. Rassias, *Stability of the generalized orthogonality functional equation*, Inner product spaces and applications, 219–240, Pitman Res. Notes Math. Ser., 376, Longman, Harlow, 1997.
6. J. Bichon, A. De Rijdt, and S. Vaes, *Ergodic coactions with large multiplicity and monoidal equivalence of quantum groups*, Comm. Math. Phys. **262** (2006), no. 3, 703–728.

¹DEPARTMENT OF MATHEMATICS, UNIVERSITY OF AAAA, BBBB 654321, CCCC, INDIA.

E-mail address: first1@afa.ac.ir; first2@afa.ac.ir

²DEPARTMENT OF PURE MATHEMATICS, FERDOWSI UNIVERSITY OF MASHHAD, P. O. BOX 1159, MASHHAD 91775, IRAN;

TUSI MATHEMATICAL RESEARCH GROUP (TMRG), MASHHAD, IRAN.

E-mail address: second@afa.ac.ir